

Water use by crops and pastures in the mid-Murrumbidgee catchment

Key points

- Agricultural production seeks both high total water use (TWU) and high water use efficiency (WUE).
- High total water use influences the environment through minimising deep drainage, and associated water logging and salinity.
- Potential or water limited yield of wheat is about 20 kg/ha/mm of seasonal water use, and for broad-leaf crops is about 15 kg/ha/mm of seasonal water use (after deducting 100 to 120 mm).
- The water limited yield of annual pasture dry matter production is about 30 kg/ha/mm (after deducting about 30 mm).
- Potentially, dryland lucerne can yield about 12 kg/ha of dry matter for each mm of annual rainfall.
- Poor WUE can indicate management problems or constraints imposed by soil, nutrition, pests or disease.

Total water use

Total water use (TWU) refers to the amount of water used by a crop or pasture. Major differences in water use are due to different rooting depths of plants. Lucerne draws water from further down the soil profile (as deep as 3–4 m) than wheat (about 1.2 m) or annual pastures based on subterranean clover (60–70 cm). Lucerne has a greater capacity to access and use water, and so minimise deep drainage. Deep drainage and its associated salinisation are minimised by high total water use by plants. However agricultural production is influenced by both the total water use and the efficiency of the plant in using water to produce grain or forage.

Water use efficiency

The efficiency of a plant to produce forage or grain is expressed as water use efficiency (WUE). WUE is a measure of the forage (biomass) or grain yield for each millimetre of water used by a pasture or crop.

Interest in Australia was spurred by the work of French and Schultz (1984a, 1984b; Figure 1) on water use efficiency in grain production by wheat.

The French and Schultz model estimates that grain yield is potentially increased by 20 kg/ha/mm of water used, after allowing for 110 mm of water ($W_{\text{threshold}}$; Figure 1).

This threshold amount of water ($W_{\text{threshold}}$) was considered by French and Schultz as water which was lost by evaporation from the soil and so not used by the crop. Later authors (Robinson and Freebairn undated) have preferred to think of it as the minimum amount of water needed before the plant will produce grain.

Other sources of unused water are in surface run off, and in deep drainage to below the rooting depth of the crop. In their data for low rainfall sites and regions (Table 1) French and Schultz had a lower $W_{\text{threshold}}$ and this would be consistent with lower evaporative loss (due to less frequent rainfall events) and to lower run off and deep drainage.




Figure 1 Relationship between grain yield of wheat and water use showing upper limits of water use (blue line, slope of 20 kg/ha/mm after allowing for 110 mm of water); data of French and Schultz 1984a (blue points); and data from five monitored crops in the Murrumbidgee catchment in 2005 (numbers).

Table 1 Estimates of water use efficiency (WUE) and threshold water ($W_{\text{threshold}}$) for grain production of wheat in South Australia and for the Murrumbidgee catchment.

Location (reference)	$W_{\text{threshold}}$ (mm)	WUE (kg/ha/mm)	Comment
South Australia (French and Schultz 1984a)	110	20	Upper limit of data (all data)
	70	20	Upper limit of data (low rainfall sites)
	170	20	Upper limit of data (high rainfall, poor soil surface conditions)
Wagga Wagga (Cornish and Murray 1989)	70	15	Average of data, dryland crops
Griffith (Steiner <i>et al</i> 1985)	126	16	Average of data, irrigated crops

Table 2 Indicative water use efficiency (WUE) and threshold water use ($W_{\text{threshold}}$) of broad leaf crops for grain production.

Crop, location (reference)	$W_{\text{threshold}}$ (mm)	WUE (kg/ha/mm)	Comment
Canola, southern NSW (Robertson and Kirkegaard 2005)	120	15	Average of data in seasons of favourable rainfall distribution
	120	11	Average (all data)
Lupins, estimate (D Lockett pers. comm.)	80	13	Approximate upper limit of data
Field pea (Siddique <i>et al</i> 2001)	100	15	Approximate upper limit of data
	100	10	Approximate average of data
Faba bean, WA estimates (Siddique <i>et al</i> 2001)	110	15	Approximate upper limit of data
	110	10	Approximate average of data

Conversely, at a site with high rainfall and a poor surface condition, where surface run off would be greater, the $W_{\text{threshold}}$ was higher (170 mm; Table 1).

Other values determined in southern NSW for wheat are given in Table 1. The $W_{\text{threshold}}$ values are in the same range as French and Schultz, but WUE is a little lower at 15–16 kg grain/ha/mm (Cornish and Murray 1989; Steiner *et al* 1985). However, these regional data are consistent with French and Schulz. They were estimates of the average values expected from reasonable agronomy and production conditions. In contrast French and Schultz described the water limited or potential maximum (upper boundary of WUE) for grain yield production.

Determining WUE will highlight inefficient use in a paddock or region. Local data from 2005 indicated reasonable WUE (Figure 1), except for *Site 4*, which is well below the water limited potential yield. WUE can be improved by identifying and correcting limitations to crop growth, such as nutrient deficiencies, diseases and weeds, soil constraints and poor agronomy such as variety choice and delayed sowing.

Water use efficiency and dry matter production

The concept of water use efficiency can be extended to vegetative production of crop and pasture. French and Schultz (1984a) indicated that wheat had a potential WUE for dry matter production (stubble and grain yield) at harvest of up to 55 kg/ha/mm, after allowing for 110mm of water loss by evaporation from the soil ($W_{\text{threshold}}$; Table 3).

Similar estimates can be made for pasture production. In Western Australia annual pastures increased their dry matter yield by 30 kg/ha/mm after allowing for 30 mm of water use (Bolger and Turner 1999; Figure 2). As this data was obtained with high input pastures, the relationship probably represents the water limited (potential) yield. The dry matter yields of low input pastures reported by Bolger and Turner (1999) were less than indicated by this relationship, and data from central NSW was below this relationship (Blumenthal and Ison 1993). Data from the Murrumbidgee catchment was near potential yield in 2006, but below potential in 2005. *Site 4* in 2005 was well below water limited potential yield.

Lucerne can produce throughout the year and its dry matter production over the year can be related to annual rainfall (Figure 3). Data from central western NSW showed that lucerne dry matter production was related to rainfall with yield increasing by 6.7 kg/ha/mm on average (Bowman *et al* 2004; Figure 3, broken line). An approximate potential of about 12 kg/ha/mm was estimated from this data (Figure 3, solid line) with no $W_{\text{threshold}}$. Again data from the Murrumbidgee catchment appears to confirm the potential yield estimate (Figure 3, numbers).

Estimating crop water use efficiency on farm

The water use efficiency concept is easiest to use in a Mediterranean climate. This is because the summers are dry, and growing season rainfall alone is a good estimate of water supply to plants. Run off and deep drainage are the unknowns, and may be unimportant in some seasons and in low rainfall regions. Crop water use is approximated by in crop rainfall (sowing to harvest).

However in the equi-seasonal environment of the Murrumbidgee catchment, where rainfall in summer is only slightly less than winter rainfall, a crop may be sown with water from summer rain already stored in the soil profile, or the crop may mature with unused water still in the soil profile. Soil moisture can be measured at the start and end of the season, or the amount of water at these times can be estimated. Farmers need some simple *rules of thumb* to assist them in these estimations.

While no simple rules can fully account for soil moisture at sowing or at harvest some guidelines are available (Robertson and Kirkegaard 2005). Working with canola in southern NSW these authors used some experimental data and simulation modelling of rainfall at Narrandera from 1904–2003 to conclude the following.

1. *Soil moisture storage at sowing* can be estimated by taking summer rainfall (post harvest to sowing), subtracting 80 mm, and then using 40% of the remainder.
2. In a similar manner, *soil moisture remaining at harvest* can be estimated as the post-anthesis rainfall (Oct–Dec), minus 50 mm, and the remainder halved. This estimate is very variable and was developed for canola, so care should be used.

An example of the calculation of WUE and water limited yield is given in the text boxes on the following page.

Table 3 Water use threshold ($W_{\text{threshold}}$) and water use efficiency (WUE) for potential dry matter production in wheat in South Australia (from French and Schultz 1984a)

Crop interval	Water use threshold ($W_{\text{threshold}}$; mm)	Water use efficiency (WUE; kg/ha/mm)
Sowing to end of tillering	40	25
Sowing to anthesis	80	60
Sowing to maturity	110	55




Figure 2 Relationship between water use and end of season dry matter production for annual pastures in WA (blue points and line; Bolger and Turner 1999), with data from the Murrumbidgee catchment in 2005 (orange numbers) and in 2006 (green numbers).




Figure 3 Relationship between annual rainfall and annual dry matter production for dryland lucerne pastures in central western NSW (blue points and lines; Bowman *et al* 2004). The broken line is the average data, and the solid line is the potential yield. Data from the Murrumbidgee catchment in 2005 (orange numbers), and in 2006 (green numbers) is presented.

Estimating WUE of a crop

Example of Site 3 in Figure 1

Sowing in early May; anthesis in early October

Rainfall

Jan–Apr	May–Dec	Oct–Dec
103 mm	457 mm	130 mm

$$\begin{aligned} A &= \text{stored water at sowing} \\ &= (103 - 80) \times 0.40 \\ &= 9 \text{ mm} \end{aligned}$$

$$\begin{aligned} B &= \text{in crop rainfall} \\ &= 457 \text{ mm} \end{aligned}$$

$$\begin{aligned} C &= \text{unused soil water at harvest} \\ &= (130 - 50) \times 0.50 \\ &= 40 \text{ mm} \end{aligned}$$

$$\begin{aligned} \text{Crop water use} &= A + B - C \\ &= 9 \text{ mm} + 457 \text{ mm} - 40 \text{ mm} \\ &= 426 \text{ mm} \end{aligned}$$

$$\text{Grain yield} = 5100 \text{ kg/ha}$$

$$\begin{aligned} \text{WUE of this crop} &= 5100 \div (426 - 110) \\ &= 16 \text{ kg/ha/mm} \end{aligned}$$

$$\begin{aligned} \text{Water limited} \\ \text{potential} &= (426 - 110) \times 20 \\ &= 6320 \text{ kg/ha} \end{aligned}$$

Estimating WUE of a crop

Your calculation

Sowing in _____; anthesis in _____

Rainfall

Jan–Sowing	Sowing–Dec	Anthesis–Dec
mm	mm	mm

$$\begin{aligned} A &= \text{stored water at sowing} \\ &= (\quad - 80) \times 0.40 \\ &= \quad \text{mm} \end{aligned}$$

$$\begin{aligned} B &= \text{in crop rainfall} \\ &= \quad \text{mm} \end{aligned}$$

$$\begin{aligned} C &= \text{unused soil water at harvest} \\ &= (\quad - 50) \times 0.50 \\ &= \quad \text{mm} \end{aligned}$$

$$\begin{aligned} \text{Crop water use} &= A + B - C \\ &= \quad + \quad - \\ &= \quad \text{mm} \end{aligned}$$

$$\text{Grain yield} = \quad \text{kg/ha}$$

$$\begin{aligned} \text{WUE of this crop} &= \quad \div (\quad - 110) \\ &= \quad \text{kg/ha/mm} \end{aligned}$$

$$\begin{aligned} \text{Water limited} \\ \text{potential} &= (\quad - 110) \times 20 \\ &= \quad \text{kg/ha} \end{aligned}$$

Limitations of the water use efficiency concept

The French and Schultz approach takes no account of the timing of rainfall. This can result in apparently low water use efficiency where, for example, severe dry conditions late in the growing season result in poor grain fill in a season with otherwise reasonable rainfall. Frost at crop flowering can also result in low apparent water use efficiency. In both these situations the crop may yield to its potential under the prevailing conditions. The farmer has done all that is possible to achieve the water limited or potential yield, but the seasonal conditions (timing of rain, frost) have limited what can be achieved.

There are also regional differences. Crops use more water under dry conditions to get the same yield as under humid conditions. Therefore, crops grown in dry areas tend to have lower WUE than in humid areas. Also in the northern wheat areas of eastern Australia soil moisture at sowing accounts for a higher proportion of water used by wheat crops than in the south. This leads to more efficient WUE because stored soil moisture is used by the crop and evaporation of stored soil moisture is negligible.

As a result of these limitations to the calculation, water use efficiency targets for water limited yield need careful use by growers. WUE is useful when comparing paddock results within a season and region. Poor WUE in a paddock, when other paddocks or growers in the same

region and season have higher WUE, probably indicates the crop yield was below what was possible; that is yield was not water limited, and there was some important management or soil based constraint on yield.

Acknowledgements

This guideline for best management practice was produced from a project conducted by Damien Doyle and Craig Muir (NSW Department of Primary Industries). The material was compiled by Dr Alison Bowman and Dr Brendan Scott, NSW Department of Primary Industries, Wagga Wagga.

Further information

www.murrumbidgee.cma.nsw.gov.au or
www.dpi.nsw.gov.au

References and further reading

Blumenthal MJ, Ison RL (1993) Water use and productivity in sub clover and murex medic swards I. Dry matter production. *Australian Journal of Agricultural Research* 44, 89-107.

Bolger TP, Turner NC (1999) Water use efficiency and water use of Mediterranean annual pastures in southern Australia. *Australian Journal of Agricultural Research* 50, 1035-1046.

Cornish PS, Murray GM (1989) Low rainfall rarely limits wheat yields in southern New South Wales. *Australian Journal of Experimental Agriculture* 29, 77-83.

French RJ, Schultz JE (1984a) Water use efficiency of wheat in a Mediterranean-type environment. I. The relationship between yield, water use and climate. *Australian Journal of Agricultural Research* 35, 743-64.

French RJ, Schultz JE (1984b) Water use efficiency of wheat in a Mediterranean-type environment. II. Some limitations to efficiency. *Australian Journal of Agricultural Research* 35, 765-75.

Robertson MJ, Kirkegaard JA (2005) Water-use efficiency of dryland canola in an equi-seasonal rainfall environment. *Australian Journal of Agricultural Research* 56, 1373-1386.

Robinson B, Freebairn D (undated) Water Use Efficiency of wheat: Principles, practices, and pitfalls. (Dept of Natural Resources, APSRU, PO Box 318 Toowoomba 4350).

Siddique KHM, Regan KL, Tennant D, Thomson BD (2001) Water use and water use efficiency of cool season grain legumes in low rainfall Mediterranean-type environments. *European Journal of Agronomy* 15, 267-280.

Steiner JL, Smith RCG, Meyer WS, Adeney JA (1985) Water use, foliage temperature and yield of irrigated wheat in south-eastern Australia. *Australian Journal of Agricultural Research* 36, 1-11.

This project has been funded through the Australian and NSW Governments' National Action Plan for Salinity and Water Quality.

This document prepared for the Murrumbidgee CMA by NSW Department of Primary Industries.



NSW DEPARTMENT OF
PRIMARY INDUSTRIES



Disclaimer

The information contained in this publication is based on knowledge and understanding at the time of writing (June 2008). However, because of advances in knowledge, users are reminded of the need to ensure that information upon which they rely is up to date and to check currency of the information with the appropriate officer of New South Wales Department of Primary Industries/Murrumbidgee Catchment Management Authority or the user's independent adviser.