

02

FarmLink Research Report 2019

Increasing the effectiveness of nitrogen fixation in pulses through improved rhizobial strains in the GRDC northern region

Report Authors

Dr Belinda Hackney, Dr Jessica Rigg, Dr Susan Orgill, Simon Flinn, Francesca Galea, Rachael Whitworth, Barry Haskins

Trial Site Location

Griffith, Condobolin, Canowindra

Introduction

Current recommendations suggest lentil and faba bean are inoculated with Group F (containing the rhizobia strain WSM1455) while field pea and vetch are inoculated with Group E (SU303). However, the current Group E is difficult to manufacture and as a result, field pea and vetch are in fact inoculated with Group F inoculant. Fortunately, the current rhizobia strain (WSM1455) used in Group F inoculant has good compatibility across all of these host legumes. However, the current Group F which was isolated from soil in Greece (pHCa 8.0) and released in 2002, exhibits a rapid decline in capacity to nodulate plants where pHCa < 6.0, with plants generally inadequately nodulated where pHCa < 5.0 (Yates *et al.* 2016; Ballard *et al.* 2019).

The aim of this project is to evaluate the potential of number of elite strains of rhizobia suitable for use across the Group E/F legume host range with a view to releasing a suitable strain that is well adapted to soil and climatic issues faced across the GRDC northern region. The release of such a strain offers potential to increase nitrogen fixation by pulses and increase the area of adaptation where they can be successfully grown.

Project Partners



Department of Primary Industries

AgGrow
AGRONOMY • RESEARCH



Funding Partners



GRDC

GRAINS RESEARCH & DEVELOPMENT CORPORATION

Project code - DAN1901-002RTX

Large areas of cropping land in central and southern NSW have soils where the capacity of the current Group F to effectively nodulate host plants is restricted. A recent survey of 150 commercial paddocks undertaken as part of this project in the GRDC northern region found that soil pH in the top 20 cm of the soil profile would likely restrict the performance of the current Group F in $\geq 40\%$ of paddocks sampled. Rhizobia with greater acid soil tolerance would likely increase capacity for nitrogen fixation throughout the regions sampled in this survey.

Both Murdoch University (Centre for Rhizobium Studies) and SARDI have been isolating rhizobia suitable for consideration as a Group F replacement from acidic soils (pHCa 4.5-5.5); Murdoch University accessions have been sourced from Italy and SARDI accessions from commercial paddocks in South Australia. Preliminary glasshouse and field studies in Western Australia identified two acid-tolerant Italian strains (WSM1455, WSM4643) which produced improved nodulation, nitrogen fixation and grain yield in field pea (Yates *et al.* 2017). Similar studies in South Australia reported improved performance in lentil and faba bean where acid tolerant Australian strains (SRDI969, SRDI970) were used compared to the current Group F (WSM1455) strain (Ballard *et al.* 2019). These results are encouraging. However, to be considered for commercial release, it is important that any potential strain has capacity to improve nodulation across the entire Group E/F host range which also includes vetch. In addition, any elite strain has to be able to be manufactured easily and to remain stable up until the point of use at sowing.

In 2019, field testing of the four elite acid-tolerant strains commenced in the Northern GRDC region with sites sown at Griffith, Condobolin and Canowindra. The objective of the three sites was to evaluate the performance of the elite strains in differing soils (Table 1) and under varying climatic conditions.

Table 1. The location, soil pH and texture of three sites used for evaluation of elite rhizobia strains for lentil, faba bean, field pea and vetch in the GRDC Northern Region in 2019.

Location	Soil pH _{Ca} (0-10 cm)	Soil texture
Griffith	4.6	Light sandy loam
Condobolin	4.8	Sandy loam (red chromosol)
Canowindra	5.1	Loam (dermosol)

For lentils, there was a significant difference in nodulation across sites and due to rhizobia strain (Figure 1). The most acidic site, Griffith, had lower nodulation than all other sites and only WSM4643 produced a significantly higher nodulation score than the current Group F strain. At Condobolin and Canowindra, both WSM1483 and WSM4643 produced significantly higher nodulation scores than the current Group F, while SRDI969 was also significantly higher than the current Group F at Canowindra. None of the strains at any site gave an overall nodulation score that is considered adequate (score ≥ 4 ; Yates *et al.* 2016).




Figure 1. The average nodulation score of 15 lentil plants at Griffith, Condobolin and Canowindra where seed was inoculated with peat slurry containing a no rhizobia (nil), the current Group F strain, or one of four experimental strains. A score of 4 is considered adequate under the system developed by Yates *et al.* (2016).

For field pea, all experimental strains produced significantly greater nodulation than the current Group F at Griffith (Figure 2). At Condobolin and Canowindra, WSM4643 produced significantly higher nodulation score than the current Group F, with WSM1483 also significantly higher at Condobolin.




Figure 2. The average nodulation score of 15 field pea plants at Griffith, Condobolin and Canowindra where seed was inoculated with peat slurry containing a no rhizobia (nil), the current Group F strain, or one of four experimental strains. A score of 4 is considered adequate under the system developed by Yates *et al.* (2016).

For vetch, WSM1483 and WSM4643 produced significantly higher nodulation than the current Group F at Griffith and Condobolin (Figure 3). None of the experimental strains produced significantly higher nodulation than the current Group F at Canowindra. SRDI970 showed poor compatibility with vetch at Condobolin and Canowindra producing very low nodulation scores. It is critical that strains considered for potential release showed strong compatibility across the potential host range.




Figure 3. The average nodulation score of 15 vetch plants at Griffith, Condobolin and Canowindra where seed was inoculated with peat slurry containing a no rhizobia (nil), the current Group F strain, or one of four experimental strains. A score of 4 is considered adequate under the system developed by Yates *et al.* (2016).

Faba beans were grown only at Canowindra and there were no experimental strains that gave nodulation scores greater than those achieved by the current Group F strain.

Samples were collected to determine nitrogen fixation and these are still undergoing processing as are samples collected to determine grain yield. However, based on results to date, it would appear that strains WSM1483 and WSM4643 offer considerable potential to increase nodulation in acidic soils and also have good compatibility with the potential host range. Research is ongoing within the Australian Inoculant Research Group to determine the manufacturability and stability characteristics of the elite strains, as this is a critical consideration in the practical delivery of elite strains to the marketplace.

Are elite strains the answer to increasing pulse production and adaptation?

The development of elite rhizobial strains is an important step forward in increasing pulse production, but it is not a silver bullet. Certainly, pulse production in central and southern NSW is likely to be constrained by acidic soils which are widespread throughout the region and our results indicate that elite strains may increase nodulation in these circumstances. The current strain of Group F was isolated from soils with a pH much higher than that encountered through much of the target growing regions of central and southern NSW and therefore strains selected from soils more comparable in pH are likely to have production and survival advantages. However, our results also show differences in host plant tolerance to soil acidity, with lentil nodulation still well below the level considered adequate even with acid tolerant rhizobia. Thus, soil acidity is potentially impacting the host plant and/or the formation of an effective symbiosis. So are tolerant host plants the answer? Potentially they are also a tool to improve adaptation and production of pulses in acid soil regions. However, what must be considered is that if the acid soil problem is not addressed, it will continue to worsen. The results of our soil survey show that a large percentage of commercial paddocks have soil pH_{Ca}<5.0. Acid tolerant rhizobia and acid tolerant host plants can only do so much and certainly once soil pH falls

below 5.0, other problems come into play including reductions in the efficiency of use of applied nutrients. Also, the impact of herbicide residues needs to be considered as these can impact the host plant, rhizobia and the host plant-rhizobia symbiosis. This project is in its early stages and further evaluation is required through time and across sites to determine the impact that elite rhizobial strains may have on potential pulse production.

References

- Ballard R, Farquharson E, Ryder M, Denton M, Rathjen J, Henry F, Brand J, Whitworth R, Haskins B, Seymour M, Yates R (2019) Fixing more N – improving the performance of rhizobial inoculants in suboptimal conditions. GRDC Updates. <https://grdc.com.au/resources-and-publications/grdc-update-papers/tab-content/grdc-update-papers/2019/02/fixing-more-n-improving-the-performance-of-rhizobial-inoculants-in-suboptimal-conditions>
- Yates, RJ, Abadidoo, Howieson, JG (2016) Field experiments with rhizobia. In. Working With Rhizobia (Eds. JG Howieson, MD Dilworth). Australian Centre for International Agricultural Research and Centre for Rhizobium Studies Murdoch University.